

Relativity in binary systems as root of quantum mechanics and space-time

W. Smilga

Isardamm 135 d, D-82538 Geretsried, Germany

e-mail: wsmilga@compuserve.com

Inspired by Bohr's dictum that "physical phenomena are observed *relative* to different experimental setups", this article investigates the notion of *relativity* in Bohr's sense, starting from a set of binary elements. The most general form of information coding within such sets requires a description by four-component states. By using Bohr's dictum as a guideline a quantum mechanical description of the set is obtained in the form of a $SO(3,2)$ based spin network.

For large (macroscopic) sub-networks a flat-space approximation of $SO(3,2)$ leads to a Poincaré symmetrical Hilbert space. The concept of a position of four-component spinors *relative* to macroscopic sub-networks then delivers the description of 'free' massive spin-1/2 particles with a Poincaré symmetrical Hilbert space.

Hence Minkowskian space-time, equipped with spin-1/2 particles, is obtained as an inherent property of a system of binary elements when individual elements are described *relative* to macroscopic sub-systems.

I. INTRODUCTION

Spin networks were introduced by R. Penrose [1] more than 30 years ago, in an attempt to describe the geometry of space-time in a purely combinatorial way. Penrose studied systems of two-component spinors which represent the simplest quantum mechanical objects. He was able to show that large systems of such spinors, generate properties of angular directions in three-dimensional space. Despite this success, the concept of $SU(2)$ based spin networks was considered not rich enough to describe also distances [2]. Penrose's interest, therefore, turned to more complex twistor objects. Nowadays spin networks are discussed as models of space-time at Planck scales [3].

Another attempt to derive space-time from "binary alternatives" was undertaken by C. F. von Weizsäcker [4,5] and co-workers. In this concept a physical particle is considered as an aggregate of 10^{40} binary alternatives, representing the knowledge about the particle. Although indications for a Poincaré symmetric space-time structure were obtained, the work remained at an experimental stage without having established a definite link to empirical particle physics.

The present article describes a new approach to derive space-time as a property of general binary systems. This goal will be achieved at 'normal' (non-Planck) scales. As a valuable guide we will repeatedly make use of Bohr's dictum that "physical phenomena are observed *relative* to different experimental setups" [6,7]. Bohr's dictum does not describe any law of nature. It rather, in a short and precise way, explains how we do physics. Because of its importance for the understanding of physics, especially at the very basic level, in this paper we will speak of *Bohr's principle of relativity* or, in short, of *Bohr's principle*.

Bohr's principle has played an important role in the interpretation of quantum mechanics. It has been used there in the sense that, the description of a quantum mechanical phenomenon must

include the experimental setup by which the phenomenon is observed. Depending on the setup an electron may appear either as a particle or as a wave. We will use Bohr's principle in the sense that "particle physics is a discipline that describes *microscopic phenomena relative to macroscopic setups*".

Our approach will start from a set of binary elements, and we will first find a way to describe the information contained in a binary element *relative* to others. We will see that in the most general case this is achieved by four-component states. Inspired again by Bohr's principle, in a next step we will introduce a quantum mechanical description of the binary elements. This leads to a spin network, made up of Dirac spinors.

Obviously, the Hilbert space of Dirac spinors is by far richer than the state space of two-component spinors that was used by Penrose. In fact, within this Hilbert space Dirac matrices generate transformations that act transitively on the state space. It is well-known that Dirac matrices thereby define an irreducible representation of the de Sitter group $SO(3,2)$. A subgroup of $SO(3,2)$ is the (homogeneous) Lorentz group $SO(3,1)$.

In large systems of Dirac spinors we can construct states, where some quantum numbers add up to large values as compared to other quantum numbers or expectation values. This then allows for a flat-space approximation by the well-known method of group contraction [8,9], which approximates the original $SO(3,2)$ symmetry by a Poincaré symmetry. The translations of the Poincaré group then generate a tangential Minkowskian space-time continuum.

This Poincaré invariant basis delivers the framework for a more detailed description of individual spinors relative to the spin network, by quantum mechanical states that carry information about their *position* within the network. These states satisfy a Dirac equation with finite mass.

Thereby we obtain a quantum mechanical system of identical, massive, non-interacting spin-1/2 particles defined in a space-time continuum with three spacelike and one timelike dimension. This system is invariant under transformations of the Poincaré group.

Since the Poincaré symmetry is only an approximation, we need to examine possible contributions of higher order terms. Such terms are defined in a natural way by the differences between $SO(3,2)$ generators and the corresponding generators of the Poincaré group. In a previous article [10] the author has shown that these terms generate interactions between particles, with characteristics of the empirical fundamental interactions, including gravitation.

II. SPINORS AND SPIN NETWORKS

Before we come to spin networks, let us consider a large system of, say 10^{80} , binary elements. Each of these elements exists in either of two *states* that we call 'up' and 'down'. Let $|u\rangle$ and $|d\rangle$ be the symbols for these states.

We can represent these states mathematically in vector form:

$$|u\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad \text{and} \quad |d\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix}. \quad (1)$$

(Here the equal-sign means ‘represented by’.) The matrix

$$\sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad (2)$$

can be used to set up an *eigenvalue equation*

$$\sigma_3|z\rangle = \zeta|z\rangle, \quad (3)$$

where $|z\rangle$ can be either $|u\rangle$ or $|d\rangle$, and ζ are the corresponding eigenvalues $+1$ and -1 , respectively. We can also define the *expectation value* of σ_3 by

$$\langle z|\sigma_3|z\rangle = \zeta, \quad (4)$$

where $\langle z|$ stands for the transposed vector of $|z\rangle$.

The matrix

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad (5)$$

applied to $|u\rangle$ or $|d\rangle$, changes $|u\rangle$ into $|d\rangle$ and $|d\rangle$ into $|u\rangle$. Since neither $|u\rangle$ nor $|d\rangle$ is preferred, we can say that σ_1 defines a discrete *symmetry transformation* on the states $|u\rangle$, $|d\rangle$.

Let us now extend the definition of a state by allowing also linear combinations of the form

$$|s\rangle = \alpha|u\rangle + \beta|d\rangle, \quad (6)$$

where α and β are complex numbers with

$$\alpha^*\alpha + \beta^*\beta = 1. \quad (7)$$

We will call such a generalized state a (two-component) *spinor*. By this step we have created a complex *vector space* with $|u\rangle$ and $|d\rangle$ serving as basis vectors. We define a scalar product of two states

$$|s_1\rangle = \alpha_1|u\rangle + \beta_1|d\rangle, \quad (8)$$

$$|s_2\rangle = \alpha_2|u\rangle + \beta_2|d\rangle \quad (9)$$

by

$$\langle s_1|s_2\rangle = \alpha_1^*\alpha_2 + \beta_1^*\beta_2. \quad (10)$$

This extends the vector space to a *Hilbert space*.

By the extension of the description of a binary element, we have not introduced any new physical degree of freedom. Still only the basis states have a physical meaning. However, the Hilbert space approach allows some mathematical operations that are not possible in the binary system. We can, for example, “close” the algebra of the matrices σ_1 and σ_3 with respect to the commutation product and obtain a third matrix σ_2 by means of

$$[\sigma_3, \sigma_1] = i\sigma_2 \quad (11)$$

in the form

$$\sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}. \quad (12)$$

It is well-known, that the matrices σ_1 , σ_2 and σ_3 , the Pauli-matrices, satisfy the commutation relations of SU(2). They serve as infinitesimal generators for SU(2) transformations on the Hilbert space of two-component spinors. SU(2) allows for continuous symmetry transformations of states within the Hilbert space, in contrast to only discrete ones between the states of the original binary elements. Thus SU(2) extends the basic discrete symmetry into the “interpolating” Hilbert space. It is, therefore, the *symmetry group* of this Hilbert space. The infinitesimal generators of SU(2) are considered as *observables* in the quantum mechanical sense.

This step has laid the basis for a quantum mechanical description of binary elements. The introduction of a Hilbert space does not change any of the “physical” properties of binary elements. It merely offers a new way of description, similar to the introduction of, for example, a potential in classical mechanics.

A *system* of spinors is then quantum mechanically described by a Hilbert space that is given by the (antisymmetrized) direct product of individual spinor states. A set of identical elements that are defined in this way and allow for the representation of a certain Lie group constitute a *spin network* [3]. Our SU(2) based system can be considered as the simplest type of a spin network.

Within such a spin network we can define sub-networks by assembling a large number of spinors. Within their Hilbert spaces we can select states with large SU(2) quantum numbers. Large quantum numbers allow for a quasi-continuous orientation of states with respect to the “direction” defined by the state of a given single spinor.

Assume that we rotate a sub-network relative to a spinor by an angle of $\pi/2$. Then by applying the corresponding SU(2) transformation to the state of the spinor we can map this rotation into the Hilbert space of the spinor. Without knowing details of the measuring process, we can conclude from symmetry arguments, that the outcome of a subsequent measurement of the spinor state *relative* to the orientation of the sub-network is non-predictable. Of course, this is familiar from the basics of quantum mechanics. This means that we have a 50:50 per cent chance to measure either an “up” or a “down” state. Therefore, the coefficients of a spinor state (6) carry information about the relation between spinor and an observer in the sense of probability amplitudes.

Coming back to Bohr’s principle, we can say that a description of a single spinor *relative* to a macroscopic sub-network, has become possible by the extension of the basic states to a vector space. Or if we formulate this statement the other way round: if we want to represent a non-discrete symmetry on a discrete system, a suitable way to achieve this is to turn to a quantum mechanical description of the system. We will come back to this point in section VII.

III. FROM SPINORS TO DE SITTER GROUP

In the last section we have taken for granted that there is only one way to code the information “up” and “down” as given by (1). However, we could have defined just as well

$$|u\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix} \quad \text{and} \quad |d\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix}. \quad (13)$$

It is a common experience in software development that programmers fail to agree on the way of how they code e.g. the outcome of a software procedure. Some may return -1 in the error case, other $+1$. Therefore, if we count errors, we cannot simply look for an outcome of -1 , but also have to consider the other way of coding. Of course, a good project management should prevent such situations arising.

Fortunately, the codings (1) and (13) represent the only ways of coding a piece of binary information in a binary system. For a single binary element the way of coding is not relevant. However, if we want to compare the information contained in several elements of a binary system, we either have to agree on one unique way of coding or we have to provide information about the way of coding with each binary element.

In the first case we can use either (1) *or* (13). In the second case the description of the information contained in a single element requires a four-component state:

$$|u_1\rangle = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \quad |d_1\rangle = \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix} \quad (14)$$

$$|u_2\rangle = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}, \quad |d_2\rangle = \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \end{pmatrix}. \quad (15)$$

These four-component objects can be understood as obtained from a direct sum of the two alternative codings of the same type of binary information. In the following we will call these objects *Dirac spinors*.

The second case has to be considered the general one, since it does not presume an agreement on the way of coding. Such an agreement, in a global sense, would be hardly compatible with relativistic causality. In other words, we cannot expect a global “project management” as far as the coding is concerned. Therefore, we consider the second case as the realistic one.

The following is based on four-component states as defined by (14) and (15).

Again we can form a Hilbert space by interpolation between the basic states. Its symmetry group is considerably larger than in the case of two-component spinors. Instead of Pauli matrices we now have 4×4 spin matrices:

$$\sigma_{ij} = \epsilon_{ijk} \begin{pmatrix} \sigma_k & 0 \\ 0 & \sigma_k \end{pmatrix}, \quad i, j, k = 1, 2, 3, \quad (16)$$

where ϵ_{ijk} is the permutation symbol. The matrix

$$\gamma^0 = \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix}, \quad (17)$$

where I is the 2×2 unit matrix, delivers an eigenvalue of $+1$, if applied to the first group of spinors (14), and -1 , if applied to the second (15). By analogy with σ_1 we can define a matrix

$$\tau = \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix}, \quad (18)$$

which interchanges the first and second group of spinors. Since none of the codings is preferred, the interchange operation again defines a symmetry transformation. Then also the operation of combined application of σ_1 and τ is a symmetry operation. It is given by the matrix

$$\gamma^1 = \begin{pmatrix} 0 & \sigma_1 \\ \sigma_1 & 0 \end{pmatrix}. \quad (19)$$

When we close the algebra of the matrices γ and σ defined so far, with respect to the commutation product, we find additional matrices

$$\gamma^k = \begin{pmatrix} 0 & \sigma_k \\ \sigma_k & 0 \end{pmatrix} \quad (20)$$

and

$$\sigma^{0k} = -\sigma^{k0} = \begin{pmatrix} 0 & i\sigma_k \\ -i\sigma_k & 0 \end{pmatrix}. \quad (21)$$

We can combine the indices 0 and k to an index $\mu = 0, \dots, 3$, and use $g_{\mu\nu} = \text{diag} (+1, -1, -1, -1)$ in the usual way to raise and lower indices.

The matrices (17) and (20) are Dirac's γ -matrices in the so-called standard or Dirac representation. They satisfy the well-known anti-commutation relations

$$\{\gamma_\mu, \gamma_\nu\} = 2g_{\mu\nu}. \quad (22)$$

and the commutation relations

$$\frac{i}{2} [\gamma_\mu, \gamma_\nu] = \sigma_{\mu\nu}. \quad (23)$$

A scalar product can be defined, as known from the treatment of the Dirac equation,

$$\langle \bar{a} | b \rangle \text{ with } \langle \bar{a} | = \langle a | \gamma^0. \quad (24)$$

The 4×4 -matrices $s_{\mu\nu}$ and $s_{\mu 4}$, built from Dirac matrices,

$$s_{\mu\nu} := \frac{1}{2}\sigma_{\mu\nu} \text{ and } s_{\mu 4} := \frac{1}{2}\gamma_\mu \quad (25)$$

form an irreducible representation of $\text{SO}(3,2)$. The proof is by verifying the commutation relations of $\text{SO}(3,2)$:

$$[s_{\mu\nu}, s_{\rho\sigma}] = -i[g_{\mu\rho}s_{\nu\sigma} - g_{\mu\sigma}s_{\nu\rho} + g_{\nu\sigma}s_{\mu\rho} - g_{\nu\rho}s_{\mu\sigma}] , \quad (26)$$

$$[s_{\mu 4}, s_{\nu 4}] = -is_{\mu\nu} , \quad (27)$$

$$[s_{\mu\nu}, s_{\rho 4}] = i[g_{\nu\rho}s_{\mu 4} - g_{\mu\rho}s_{\nu 4}]. \quad (28)$$

The transformations of $\text{SO}(3,2)$ extend the discrete symmetry operations of the general binary system, to the interpolating Hilbert space of Dirac spinors. Therefore, $\text{SO}(3,2)$ is the basic symmetry group of Dirac spinors. A subgroup of $\text{SO}(3,2)$ is the homogeneous Lorentz group $\text{SO}(3,1)$ with the commutation relations (26). The common subgroup $\text{SO}(3)$ reflects the $\text{SU}(2)$ symmetry of the two-component spinor parts of the Dirac spinor.

A *system* of Dirac spinors is described by the direct product of individual Hilbert spaces. The operators

$$S_{\mu 4} = \sum s_{\mu 4}, \quad S_{\mu\nu} = \sum s_{\mu\nu} \quad (29)$$

then generate $\text{SO}(3,2)$ symmetry transformations and are considered as observables.

IV. POINCARÉ GROUP AND SPACE-TIME

Consider now a *large* sub-network within a spin network of Dirac spinors. From these spinors we can construct states with large quantum numbers. These quantum numbers can then be regarded as quasi-continuous. Assume that we have constructed these states in such a way that the expectation values of the operators $S_{\mu 4}$ are large compared to those of $S_{\mu\nu}$.

Then we can apply group contraction [8,9] to this sector of the Hilbert space, and thus obtain the Poincaré group $\text{P}(3,1)$ as an approximate symmetry group. The operators $S_{\mu 4}$ are then approximated by translation operators

$$S_{\mu 4} \rightarrow P_\mu \quad (30)$$

with

$$[P_\mu, P_\nu] = 0 \quad (31)$$

and

$$[S_{\mu\nu}, P_\rho] = i[g_{\nu\rho}P_\mu - g_{\mu\rho}P_\nu]. \quad (32)$$

The quasi-continuous spectrum of quantum numbers is now replaced by the continuous one of the operators P_μ .

Eigenstates of P_μ can be used to form new states $|X\rangle$ that are *localized* in spacelike directions

$$|X\rangle = (2\pi)^{-3/2} \int_{-\infty}^{\infty} d^3P e^{ix^\mu P_\mu} |P\rangle. \quad (33)$$

The operators P_μ generate translations when applied to the states $|X\rangle$

$$e^{ia^\mu P_\mu} |X\rangle = |X + a\rangle. \quad (34)$$

The parameter space of the offsets a then defines a 4-dimensional *space-time* continuum with Minkowskian metric.

This means that we have derived space-time as a property of large sub-networks in the form of a tangential space-time continuum. Tangential means: it is a flat-space approximation, obtained by group contraction, valid for the neighbourhood of a given point in space-time. In section VI we will indicate how more accurate approximations can be obtained.

V. SPINORS IN SPACE-TIME

After having introduced the concept of space-time for large or “macroscopic” sub-networks, Bohr’s principle motivates us to try to find a description of a single spinor relative to the positions of sub-networks.

In the following, we will use the term “macrosystem” for a macroscopic sub-network that is described by the Poincaré invariant approximation.

Let $|P\rangle$ be the state of a macrosystem with momentum P , and let $|s\rangle$ be the state of a separate Dirac spinor that is not part of the macrosystem. Then the combined state is obtained from the direct product $|P\rangle|s\rangle$. For sake of clarity we add the operators corresponding to $S_{\mu 4}$

$$(P_\mu + s_{\mu 4})|P\rangle|s\rangle. \quad (35)$$

Since the Dirac spinor can also be considered as part of an extended macrosystem, there exists an alternative representation in the form

$$P_\mu|P'\rangle = P_\mu|P + p_s\rangle, \quad (36)$$

which describes the macrosystem including the spinor. In the Poincaré invariant approximation, on which we have based this consideration, P' can differ from P only by an incremental 4-momentum p_s , which may depend on the “internal” quantum numbers of $|s\rangle$. The momentum p_s can be considered as the effect on the macrosystem that is caused by the inclusion of the spinor.

Let us now decide that in the following the momentum operator P_μ shall act only on the original macrosystem (as in (35)). To compensate for this decision we introduce a second momentum operator p_μ that acts on the momentum p_s . Suppose that the momentum state on the rhs of (36) is given in the form

$$e^{ix_\mu(P_M^\mu + p_s^\mu)} = e^{ix_\mu P_M^\mu} e^{ix_\mu p_s^\mu}, \quad (37)$$

where P_M^μ and p_s^μ are quantum numbers of macrosystem and spinor, respectively. Then we can split (36) in the following way

$$(P_\mu + p_\mu)|P\rangle|p_s\rangle. \quad (38)$$

When we compare (35) with (38) we find the following correspondence

$$s_{\mu 4}|s\rangle \Longleftrightarrow p_\mu|p_s\rangle. \quad (39)$$

We can understand this in the following way: In combination with the state of a macrosystem, the action of $s_{\mu 4}$ on $|s\rangle$ is simulated by the action of p_μ on an auxiliary state $|p_s\rangle$.

The dependency on s makes $|p_s\rangle$ again a four-component state. Therefore, we can map this state into the Hilbert space of Dirac spinors. Let $|\psi_s(p)\rangle$ be the image of $|p_s\rangle$. Then $|\psi_s(p)\rangle$ is a Dirac spinor defined as a superposition of the four basic states with coefficients depending on p_s . This mapping has to be covariant with respect to transformations of the Poincaré group. Therefore, to determine the coefficients, we need a rule that is invariant with respect to P(3,1). The only invariant rule that can be formed from the characteristic operators p_μ and $s_{\mu 4}$ of relation (39) and that is linear in both operators can be written, with $s_{\mu 4} = \frac{1}{2}\gamma_\mu$, as

$$(\gamma^\mu p_\mu - m)|\psi_s(p)\rangle = 0. \quad (40)$$

This is the Dirac equation with a mass of m . So the Dirac equation appears here as a relation that links representation (38) to (35).

A value for the mass m is obtained by the following consideration. The operator $p^\mu p_\mu$ “simulates” the operator $\frac{1}{4}\gamma^\mu\gamma_\mu$, which evaluates to the unit matrix. Therefore, a reasonable estimate for m is the value of 1 (measured in units of quantum numbers).

By this manipulation we have formally given a single spinor a momentum in relation to an arbitrary macrosystem. This does not mean that the spinor has obtained additional degrees of freedom. There are still only four independent states in the Hilbert space of a spinor. The momentum information is instead rather coded into the coefficients of the spinor states. These coefficients are essentially phase factors and reflect the way how the spinor is coupled to a macrosystem.

Because we have “borrowed” the momentum p_s from the macrosystem, the transformation properties of p_s with respect to the Poincaré group are identical to that of P . From the momentum states $|\psi_s(p)\rangle$ we again can form states that are formally localized in spacelike directions. Then the translations of the Poincaré group will move these localized states in space-time. Such a state may “start” from a macrosystem A, and then “arrive” at the position of another macrosystem B. Again we can form a direct product with macrosystem B, and obtain the effect which this spinor causes to B. However, between A and B the position of the spinor is not defined. This does not cause any problems, but reflects the fact that we do not have information about the position of the spinor between A and B. But let us insert an experimental setup C in between A and B, to observe the spinor on its way from A to B. Then we can form a product state of the spinor and the setup C, and verify that it was in fact on its way from A to B, but was caught at C.

Thus we have arrived at a logically consistent description of a *spinor in space-time*. The apparent space-time properties of the spinor do not reflect degrees of freedom of the spinor itself, but stand for the *relation* of the spinor to macrosystems. If there were no other spinors than the one under consideration, we would not have been able to introduce any space-time properties.

VI. INTERACTIONS

The flat-space approximation obtained by group contraction cannot be the final answer to the kinematical properties of a $SO(3,2)$ based spin network. The following considerations will show why.

Let us come back to a macroscopic sub-network, described by momentum P , and a single spinor s , which form a common state $|P, s\rangle$. In flat-space approximation, translations of this state are generated by the momentum operator P_μ . In an exact $SO(3,2)$ description, we have instead a state $|S, s\rangle$, and the operator

$$S_{\mu 4} + s_{\mu 4} \quad (41)$$

generates symmetry transformations that correspond to the translations in flat-space approximation. In performing the contraction limit, where the contributions of $S_{\mu 4}$ are assumed to grow over all limits, the contributions of $s_{\mu 4}$ to the translations usually are ignored relative to those of $S_{\mu 4}$.

But in the last section the situation is different. Although the momentum operator p_μ has been derived from the momentum P_μ of the macroscopic system, the contribution of p_μ can no longer be considered large with respect to that of $s_{\mu 4}$. In fact, since p_μ stands for the effect of $s_{\mu 4}$ on the macrosystem, both operators have the same magnitude. This requires an improvement of the approach in the last section by taking into account $SO(3,2)$ specific elements.

We may start from $SO(3,2)$ states $|S\rangle|s\rangle$ that are given by the direct product of a quasi-continuous representation of a macroscopic sub-network and a spinor representation. By reduction with respect to $SO(3,2)$ we obtain

$$|S\rangle|s\rangle \rightarrow \sum_s |S_s, s\rangle = \sum_s |S + d_s, s\rangle. \quad (42)$$

The target is then to factorize the state on rhs of (42) such that

$$\sum_s |S + d_s, s\rangle = |S\rangle \sum_s |d_s, s\rangle. \quad (43)$$

Then again the sub-network $|S\rangle$ can be separated and we obtain a $SO(3,2)$ description of a spinor relative to a sub-network by states

$$|d_s, s\rangle. \quad (44)$$

This description has the form of a direct product of a quasi-continuous representation and a spinor representation. Whether or not this target can generally be reached is not known.

However, an approach starting from exact $SO(3,2)$ states, as just sketched, is not what is really required, for the following reason: Experimental setups are macrosystems in a Poincaré invariant

environment. If we try to apply corrections to the kinematics which are required by $SO(3,2)$ symmetry, we, nevertheless, have to measure the consequences of these corrections *relative to a Poincaré invariant basis*. Therefore, we need a theoretical approach that is adaptive to this situation.

This demand can be satisfied by a perturbational approach that treats the differences between the generators of $SO(3,2)$ and $P(3,1)$, for example the term $s_{\mu 4}$, as perturbations to a Poincaré invariant basic system. Whether or not the perturbation algorithm converges to separable states in the sense of (43) is of secondary interest. Of physical interest are rather, the consequences of corrections caused by $SO(3,2)$ symmetry. These corrections are obtained by employing $SO(3,2)$ operators, but do not require a knowledge of the exact $SO(3,2)$ states.

Such a perturbative approach has been described in a recent article by the author [10]. The article is based on a spin network with $SO(3,2)$ representations in the sense of (44). It treats the spin network by perturbation methods relative to a Poincaré invariant basis. This leads to the identification of four types of perturbation terms. It is shown that these terms give rise to interactions between particles with characteristics of the four empirical interactions: electromagnetic, weak, chromodynamic and gravitational.

VII. BOHR'S PRINCIPLE AND QUANTUM MECHANICS

We have intentionally started from an set of binary elements rather than from a spin network. We then have introduced a Hilbert space that interpolates between the states of the binary elements, and have obtained $SU(2)$ and $SO(3,2)$ based spin networks.

Macroscopic sub-networks are described by quasi-continuous representations and allow, for example, quasi-continuous rotations in space. According to Bohr's principle, we had to find a way to describe the discrete system of a single spinor *relative to a macroscopic* setup that can be randomly rotated in space. Obviously, a practicable way is to use a description that interpolates between the basic states of the microscopic system.

So our answer to the question "Why quantum mechanics?" is simply this: Quantum mechanics is a way to fulfil Bohr's principle in the microscopic domain.

We have introduced quantum mechanics at the binary level in a simple and rather formal way. If we had not known quantum mechanics before, we could have hardly anticipated the extreme power of quantum mechanics in general. Under this perspective, the success of quantum mechanics can be ascribed mainly to the following properties: Firstly, the quantum mechanical calculus can be seamlessly extended from binary systems to complex composed systems, by forming direct products of Hilbert spaces of subsystems. Secondly, quantum mechanics keeps track of probabilities, and composed systems inherit this property from their parts. This delivers information of statistical nature in situations where no deterministic information is available. Lastly, in "classical" limits probabilistic properties become deterministic and may even manifest themselves in form of additional independent degrees of freedom. An example in this context are momentum and position. They become independent degrees of freedom, although position, at the quantum mechanical level, is defined only by superposition of momentum states.

So the introduction of an interpolating Hilbert space appears as a mathematical trick that in the end is more than justified by practical results.

VIII. CONCLUSION

Bohr's principle has guided us from a most general binary structure, to physical concepts that in the end, allows us to describe observations relative to experimental setups. These concepts include the basics of quantum mechanics, space-time, particles and interaction.

It is very satisfactory, that as a consequence of this approach a Minkowskian space-time continuum is obtained. Thus space-time appears as a derived phenomenon, that stands for a way of describing subsystems within binary systems relative to each other.

Note that we have obtained space-time as a property of *macroscopic* systems. This is in contrast to (so far unsuccessful) attempts to derive space-time from discrete structures at Planck scales. Nevertheless, our approach is also basically a discrete one, but discrete at microscopic scales rather than at Planck scales.

The symmetry operations that have been made available by the basic symmetry group $SO(3,2)$ deliver the basis for rearrangements of relations and especially for different reference frames. Thus Bohr's principle of relativity can be considered as the root of relativistic principles, including special and general relativity, and also of dynamical processes within spin networks.

But first of all Bohr's principle has guided us to a description of *microscopic phenomena relative to macroscopic setups* by interpolating Hilbert spaces. Therefore, quantum mechanics can be understood as a necessary logical result of the application of Bohr's principle to the microscopic domain.

Our results exhibit a general binary structure as a basis, from which essential aspects of particle physics can be derived. Is then a binary system the ultimate answer to our search for basic structures in nature? Does it describe some kind of a pre-physical aspect of nature, or is it rather a logical consequence of our continuous search for more and more elementary structures? We know from information technology that we arrive at binary structures, when we divide information into smaller and smaller pieces. Do we come across the same situation in particle physics?

To quote another dictum of Bohr's: "Physics is not about reality, but about what we can say about reality". In this sense, binary systems can be regarded as a general means to represent our knowledge about nature at the most elementary level. But then the binary system itself is not part of the knowledge. It is merely an unspecific logical scheme, comparable to the binary system in computer technology, that is used to represent information but is not part of the information. Since binary information cannot be divided further, there cannot be any knowledge beyond what can be coded into the binary structure.

Therefore, we cannot say that nature has the structure of a binary system. The binary system rather reflects our way of doing science. We typically practise science by dividing knowledge into smaller and smaller parts. At least we can say that our knowledge about nature at microscopic scales can be represented within a discrete, binary logical scheme. At the same time we have to accept the

fact that the binary structure sets a logical limit to further refining of knowledge. Also: there are no “laws of nature” at the binary level other than the laws of mathematical logic.

If this is the “final answer”, then quantum mechanics, space-time, particles and interactions are inevitable logical consequences, not of fundamental laws of nature, but of the way in which we do physics. This way is determined by two of our behaviour patterns: firstly, by our habit to divide information into smaller and smaller parts, and, secondly, by the application of Bohr’s principle of relativity to observe and describe relations between these parts.

This view illustrates another famous dictum of Bohr’s: “It is the task of science to reduce deep truths to trivialities”.

-
- [1] R. Penrose, “Angular momentum: an approach to combinatorial space-time”, in: *Quantum Theory and Beyond*, ed. Ted Bastin (Cambridge University Press, Cambridge, 1971).
 - [2] R. Penrose, W. Rindler, *Spinors and space-time*, p. 43, (Cambridge University Press, Cambridge, 1988)
 - [3] J. C. Baez, “An introduction to spin foam models of quantum gravity and BF theory”, in: *Geometry and Quantum Physics*, eds. H. Gausterer and H. Grosse, (Springer, Berlin, 2000). Available as gr-qc/9905087.
 - [4] C. F. von Weizsäcker, “Binary Alternatives and Space-Time Structure”, in: *Proceedings, Quantum Theory and the Structure of Time and Space, Vol.2 (In Memoriam Werner Heisenberg)* (Munich, 1977)
 - [5] C. F. von Weizsäcker, “Urs, Particles, Fields”, in: *Proceedings, Quantum Theory and the Structure of Time and Space, Vol.5* (Munich, 1983)
 - [6] As quoted in M. Jammer, *The Philosophy of Quantum Mechanics* (Wiley, 1974)
 - [7] I am indebted to the following article for drawing my attention to Bohr’s dictum:
Djordje Minic, Chia-Hsiung Tze, “A general theory of quantum relativity”, hep-th/0309239
 - [8] E. Inönü and E. P. Wigner, *Proc. Nat. Acad. Sci. USA* **39**, 510 (1953).
 - [9] F. Gürsey in: *Group Theoretical Concepts and Methods in Elementary Particle Physics*, ed by Gürsey (Gordon and Breach, 1964).
 - [10] W. Smilga, “Spin foams, causal links and geometry-induced interaction”, hep-th/0403137